

# Foot Injury

## Definition

- Minor injury to the foot.
- Your child can walk, but with mild pain.
- If unsure, contact your child's doctor.

## Health Information

### Types of Foot Injuries

- Sprains - stretches and tears of foot ligaments.
- Strains - stretches and tears of foot muscles (a pulled muscle).
- Muscle bruise from a direct blow.
- Bone bruise from a direct blow.
- Fractures (broken bones).
- Dislocations (bone out of joint).

### Symptoms of Foot Injuries

- The main symptom is pain and tenderness of the injured area.
- The pain is made worse with foot movement and walking.
- Swelling and bruising may also occur there.

### Prevention

- Supervise young children carefully.
- Teach older children about high-risk behaviors.
- Wear closed-toe supportive shoes (like sneakers) when physically active.

## Care Advice

### 1. Key Points

- Most foot injuries can be treated at home.
- Here is some care advice that should help.

### 2. First Aid for Foot Injuries

- Apply a compression wrap (elastic bandage) right away to limit bleeding and swelling. Reason: The more the initial bleeding and swelling, the longer the recovery.
- Apply crushed ice in a plastic bag or a cold pack for 20 minutes.
- Elevate and rest the injured foot for 24 hours.
- After 24 hours of rest, allow any activity that doesn't cause pain.

### 3. Cold Pack for Initial Pain

- For pain, swelling or bruising with any foot injury, use a cold pack. You can also use ice wrapped in a wet cloth.
- Put it on the area for 20 minutes.
- Repeat for 20 minutes every hour for the first 4 hours, then only as needed.
- Reason: Helps with the pain and helps stop any bleeding.
- Caution: Avoid frostbite.

#### 4. **Heat for Pain After 48 Hours**

- If pain lasts over 2 days, put heat on the sore area.
- Use a heat pack, heating pad or warm wet washcloth.
- Do this for 10 minutes, as needed. Reason: increases blood flow and speeds healing.
- Caution: Avoid burns.

#### 5. **Pain Medicine**

- For pain relief, give acetaminophen every 4 hours or ibuprofen every 6 hours as needed (See Dosage table).
- Ibuprofen may be more effective for this type of pain.
- Also continue compression with the elastic bandage as long as it helps the pain.

#### 6. **Activity and Movement During Recovery**

- Rest the injured part for the first 24 hours.
- Then start moving the foot as tolerated. Do this at least 3 times per day.
- Reason: This helps prevent tight muscles and joints. It helps maintain range of motion.
- As pain improves, slowly increase activity. Allow any activity that does not cause pain.
- Here's a good rule: If it HURTS, don't do it.

#### 7. **What to Expect:**

- Pain and swelling usually peak on day 2 or 3.
- Swelling is usually gone by 7 days.
- Pain may take 2 weeks to completely resolve.
- Protect the area from further injury until healed.

### **Call Your Doctor If**

- Pain not improved after 3 days
- Pain or swelling lasts over 2 weeks
- Can't walk
- You think your child needs to be seen
- You have other questions or concerns

#### **Pediatric Care Advice**

**Author:** Barton Schmitt MD, FAAP

**Copyright 2000-2023** Schmitt Pediatric Guidelines LLC

**Disclaimer:** This health information is for educational purposes only. You the reader assume full responsibility for how you choose to use it. The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. Listing of any resources does not imply an endorsement.